



RED HEN

SE17ENTEEN
17% Protein Premium Layer



DEWORMER

Laucke Mills



www.lauckemills.com.au
AUSTRALIAN OWNED AND MADE
Since 1899



RED HEN

DEWORMER

RED HEN DEWORMER CONTAINS 60PPM FLUBENDAZOLE

- Highly effective dewormer
- No withholding period for eggs
- Ease of dosing
- Use for worming programs for the control of the most common worm species found in chickens, including tapeworms. *Ascaridia galli*, *Heterakis gallinarum*, *Capillaria spp.*, *Amidostomum Anseris*, *Trichostrongylus tenuis* and *Raillietina sp.*
- Flubendazole's efficacy against all life stages of the worm (adult, larvae and eggs), means the beneficial effect on the flock is long-lasting. There is no need to discard eggs during or after treatment periods.
- Signs of infestation can be difficult to identify. Rate of lay, egg size, yolk colour and egg shell quality can all be affected by worm burdens.

DOSAGE

Feed for 7 consecutive days.

Use this feed within the best before date of 6 months of manufacture.

Remove all other feed and ensure birds have access to feed *ad lib.* to enable consumption of the required dose.

TREATMENT PERIODS

Treatment periods can be covered by 2 approaches.

A routine schedule: 2-4 times per year
Or

Faecal floatation (via diagnostic labs or vet clinics) to detect the severity of the burden on the bird and if deworming is required.

INFECTION PRESSURE*	FREQUENCY
High	3-4 weeks
Medium	6-8 weeks
Low	8-10 weeks

*Infection Pressure is affected by stocking density and environmental contamination. Environmental hygiene management is essential for optimum worm control.

WITHHOLDING PERIODS

MEAT: Remove medicated feed 7 days before slaughter for human consumption.

EGGS: Nil

CONTRAINDICATIONS

The use of Flubendazole is contraindicated in pigeons and parrots.

MADE FROM



RED HEN

17



Laucke Mills, 1519 Sturt Highway, Daveyston
PO Box 11, Greenock, South Australia, 5360
Phone: (08) 8562 8140 email: info@lauckemills.com.au